- (iii) Fill in the blanks: 1×5=5
  - (A) D-glucose is an epimer of
  - (B) Ketoses have less number of than aldoses.
  - (C) Mild oxidation of glucose gives
  - (D) \_\_\_\_\_ is present mostly as furanose.
  - (E) The common form of glucose as represented by Haworth projection is known as
- (h) (a) Why is ESR spectrum recorded in derivative mode?
  - (b) How many ESR lines are observed in methyl radical? Explain. 2
  - (c) In which region of the δ scale usually aromatic hydrogens absorb in a <sup>1</sup>H NMR spectrum and why?
  - (d) How would you expect the <sup>1</sup>H NMR spectrum of ethanol to vary when it is recorded as—
    - (i) pure ethanol;
    - (ii) ethanol in presence of small amount of water?

total number of printed pages -12

3 (Sem-6/CBCS) CHE HC 2

## 2022

## CHEMISTRY

(Honours)

Paper: CHE/HC/6026

(Organic Chemistry-V)

Full Marks: 60

Time: Three hours

The figures in the margin indicate full marks for the questions.

Answer the following questions : (any seven)  $1 \times 7 = 7$ 

- (a) Give an example of edible dye.
- (b) Which one of the following is most reactive for amonic polymerization?
  - (1)  $CH_2 = CH NO_2$
  - $(u) \quad CH_2 = CH CH_3$

- (iii)  $CH_2$  CH  $C_1H_1$
- (iv)  $CH_3 C = CII$ ,  $CH_3$
- (c) Which of the following is laevorotatory?
  - (i) Glucose
  - (ii) Fructose
  - (m) Sucrose
  - (iv) Cellulose
- (d) Fill in the blank

The auxochrome group in the pieric acid is

- (c) The electronic transition, which requires maximum energy is
  - (1)  $\sigma \sigma^*$
  - (ii)  $\pi \to \pi^*$
  - (iii)  $n \to \pi^*$
  - (iv)  $n \rightarrow \sigma^*$

- (f) Which of the following compounds absorb UV radiation?
  - (i) Heptane
  - (ii) Benzene
  - (iii) Butadiene
  - (iv) Acetone
- (g) Which of the following compounds does not show mutarotation?
  - (i) Glucose
  - (ii) Fructose
  - (iii) Maltose
  - (iv) Sucrose
- (h) How many stereoisomers should an aldohexose have?
- (i) Ribose and xylose are
  - (i) epimers
  - (ii) anomers
  - (ni) disaccharide
  - (iv) optically inactive
- (j) What are the constituents of starch?

## 2. Give answer of the following: (any four) $2\times4=8$

- (a) Write the expected products by showing the reaction of hydrolysis of lactose.
- (b) Glucosides neither give positive test with Fehling solution or Tollen's reagent nor undergo mutarotation. Explain.
- (c) Indicate the mechanism, cationic, anionic or free radical—by which the following monomers will undergo polymenzation:

(1) 
$$CII_2 = C < \frac{CH_3}{CH_3}$$

(ii) 
$$CF_2 = CF_2$$

(iii) 
$$CH_2 = CH - OCOCH_3$$

(iv) 
$$CH_2 = C \frac{CN}{CN}$$

(d) Give the method of preparation and uses of PVC and neoprene.

- (e) How do you explain the greater stability  $\beta D(+)$ -glucopyranose?
- (f) Why is the  $\lambda_{max}$  for the diene (I) low than diene (II).

$$(I) \qquad (II)$$

- (g) "Though azobenzene is a coloured compound it is not used as a dye." Explain why.
- (h) Fill in the blanks:
  - (i) Amylose is a \_\_\_\_\_ polymer of
  - (ii) Amylopectin is a \_\_\_\_\_ polymer of \_ \_\_\_.
- 3. Answer **any three** of the following: 5×3=15
  - (a) (i) Draw the cyclic anomeric forms of D-fructose.
    - (ii) Give the mechanism for hydrolysis of glycoside under acidic condition. 1+4=5

- (b) Explain the following:  $2\frac{1}{2} \times 2=5$ 
  - (i) Chemical shift
  - (ii) Spin-spin coupling
- (c) (i) Differentiate thermoplastic and thermosetting polymers.
  - (ii) Give the mechanism of acid catalyzed formation of phenol-formaldehyde resin.

 $2\frac{1}{2} \times 2 = 5$ 

- (d) How many proton signals would be expected in NMR spectra of each of the following compounds? 2½×2=5
  - $\mu$  =  $CCH_3CH_2CH_2OH$
  - (ii)  $CH_3 = O CH_2 CH_3$
- (c) Differentiate the following by giving one example of each:

 $2\frac{1}{2} \times 2 = 5$ 

- (i) Reducing sugar and non-reducing sugar
- (ii) Sugar and non-sugar

- (f) Find out the correct answer of the following: 1×5=5
  - (i) Glucose cannot be clarified as (hexose, an oligosaccharide, an aldose, a monosaccharide)
  - (ii) The monosacchande obtained by hydrolysis of starch is(D-glucose, maltose, D galactose, D ribose)
  - (m) The product which is not derived from cellulose is, (rayon, insulin, gun cotton, paper)
  - (iv) Carbohydrates are stored in the body as (sugars, starch, glucose, glycogen)
  - (v) Hydrolytic conversion of sucrose into glucose and fructose is called (induction, insertion, inversion, inhibition)
- (g) (i) A very strong characteristic absorption for -C = C stretching vibration is observed for cv.? butene but not for trans-2-butene Explain briefly. 2

- (n) A compound A having molecular formula  $C_3H_6O$  gave the following IR spectral data:

  17.00cm 1 and 2820cm-1 (doublet) and 17.30cm-1 (singlet).

  Deduce the structure of the compound A and also explain the spectral data.
- (h) (i) What is a leuco base? How can it be converted into a dye?
  - (ii) How will you synthesize alizarin from anthraquinone?

 $2\frac{1}{2} \times 2 = 5$ 

- 4. Answer **any three** of the following:

  10×3=30
  - (a) (i) What is Ziegler-Natta polymerization? Discuss its special importance in the synthesis of addition polymers.

1+4=5

(11) What is Nylon-66?


Rubber

2

(iii) Write the structures of monomer unit for the following polymers:

Polyvinyl chloride, Teflon and

- (b) (i) A pleasant smelling liquid having molecular formula  $C_9H_{10}O_2$  shows three singlets in the NMR spectrophotometry at  $\delta$  7.31 (5H), 5.08 (2H) and 2.06 (3H) and an IR peak at 1730cm<sup>-1</sup> but none near 3350cm<sup>-1</sup>. Identify the compound.
  - (ii) What kind of transition of the compound  $CH_3OCH_3$  gives rise to the 185nm absorption?
  - (iii) Which one of the following would be expected to absorb light of longest and shortest wavelength and why?



- Str.(A)
- Str.(B)
- Str.(C)
- (c) (i) Explain the following:  $2\times2=4$ 
  - (a) H bonding raises the wavelength of absorption.
  - (b) I effect raises the wave number of absorption.

 $CH_2$ 

- (ii) How will you distinguish the following by spectroscopy? 3×2=6
  - (A) Salicyclic acid and p-hydroxybenzoic acid (by IR).
  - (B)  $ClCH_2CH_2Cl$  and  $CH_3CHCl_2$  (by  $^1H$  NMR)
- (d) (i) Give the structural formula of the following: 2×3=6
  - (a) Fluorescein
  - (b) Congo red
  - (c) Methyl Orange
  - (ii) What Chromophore is group present in— 1×2=2
    - (a) fluoroscein in alkaline medium;
    - (b) malachite green?
  - (iii) Which one of the following is highly coloured?

- (w) What is Witt's theory of colour and constitution of dye?
- (c) (i) Give the concept of poly-dispersion in polymers.
  - (ii) How will you synthesize polystyrene from benzene? 3
  - (iii) State the differences between addition and condensation polymerization.
  - (iv) Give reasons why PVC is soft and flexible whereas bakelite is hard and brittle.
- (f) Write notes on the following:  $2\times5=10$ 
  - (i) Co-polymerisation
  - (ii) Rubber
  - (iii) Configuration of polymer chains
  - (w) Polymer classification
  - (v) Electrically conducting polymers
- (g) (i) Explain why the polysaccharide do not mutarotate. 2
  - (11) Give the structures of sucrose, lactose and maltose.